




Testing general relativity with black-hole binary observations: a manifesto

Michele Vallisneri
Jet Propulsion Laboratory, Caltech




- Detecting GWs offered a profound confirmation of general relativity
- Using GW observations to test GR to increasing precision, we hope to obtain clues of new physics, and ultimately prove a more general theory



Testing GR with GW observations: a **manifesto**

- **Consistency:** a useful sanity check, but hard to interpret statistically. P values are possible with much work. But would we ever believe an inconsistent result?


the graphical model of parameter inference





GW150914 tests of “consistency”: partial waveforms (1)




GW150914 tests of “consistency”: partial waveforms (1)





GW150914 tests of “consistency”: partial waveforms (2)



GW150914 tests of “consistency”: partial waveforms (2)



GW150914 test of “consistency”: full-waveform residual



SNR in coherent burst analysis
of data residual after subtracting
best-fit GW150914 waveform


$$\text{SNR}_{\text{res}}^2 = \frac{1 - \text{FF}^2}{\text{FF}^2} \text{SNR}_{\text{det}}^2$$

Fitting Factor: parameter-maximized
waveform overlap

$$\text{SNR}_{\text{res}} \leq 7.3 \Rightarrow \text{FF} \geq 0.96$$

(for violations not absorbed
by physical parameters)

GW150914 test of “consistency”: full-waveform residual



$$\text{SNR}_{\text{res}}^2 = \frac{1 - \text{FF}^2}{\text{FF}^2} \text{SNR}_{\text{det}}^2$$

Fitting Factor: parameter-maximized waveform overlap



$$\text{SNR}_{\text{res}} \leq 7.3 \Rightarrow \text{FF} \geq 0.96$$

(for violations not absorbed by physical parameters)

Testing GR with GW observations: a **manifesto**

- **Consistency:** useful sanity checks, but hard to interpret statistically. P values are possible with much work. But would we ever believe an inconsistent result?
- **Parametric tests:** constraints on GR “constants” (PN coefficients, graviton mass) are useful proxies for increasing resolving power, but again hard to interpret. Apparent violations may focus our search for new physics.

GW150914 parametric test of GW propagation: graviton mass




$$h(f) = \frac{1}{D} \frac{\mathcal{A}}{\sqrt{\dot{F}}} f^{2/3} e^{i\Psi(f)}$$

$$\Psi(f) = \sum_i [\psi_i + \psi_{il} \log f] f^{(i-5)/3} + \Phi^{\text{MR}}[\beta_i, \alpha_i]$$

$$\frac{v_g^2}{c^2} = 1 - \frac{m_g^2 c^4}{E^2}$$

$$\delta\Psi(f) = \frac{\pi D c}{\lambda_g^2 (1+z) f}$$

GW150914 parametric test of GW propagation: graviton mass





$$h(f) = \frac{1}{D} \frac{\mathcal{A}}{\sqrt{\dot{F}}} f^{2/3} e^{i\Psi(f)}$$

$$\Psi(f) = \sum_i [\psi_i + \psi_{il} \log f] f^{(i-5)/3} + \Phi^{\text{MR}}[\beta_i, \alpha_i]$$

$$\frac{v_g^2}{c^2} = 1 - \frac{m_g^2 c^4}{E^2}$$

$$\delta\Psi(f) = \frac{\pi D c}{\lambda_g^2 (1+z) f}$$


GW150914 test of generation and binary dynamics: frequency-domain PN coefficients



$$h(f) = \frac{1}{D} \frac{\mathcal{A}}{\sqrt{\dot{F}}} f^{2/3} e^{i\Psi(f)}$$

$$\Psi(f) = \sum_i [\psi_i + \psi_{il} \log f] f^{(i-5)/3} + \Phi^{\text{MR}}[\beta_i, \alpha_i]$$

GW150914 test of generation and binary dynamics: frequency-domain PN coefficients





$$h(f) = \frac{1}{D} \frac{\mathcal{A}}{\sqrt{\dot{F}}} f^{2/3} e^{i\Psi(f)}$$

$$\Psi(f) = \sum_i [\psi_i + \psi_{il} \log f] f^{(i-5)/3} + \Phi^{\text{MR}}[\beta_i, \alpha_i]$$

For comparison: pulsar-timing tests of GR concern physical parameters, but also have weak interpretations

PSR B1913+16
[Weisberg 2003]



PSR J0737-3039
[Kramer 2016]

$$\begin{aligned}\dot{P}_b &= -\frac{192\pi}{5} \left(\frac{P_b}{2\pi}\right)^{-5/3} \left(1 + \frac{73}{24}e^2 + \frac{37}{96}e^4\right) (1-e^2)^{-7/2} T_\odot^{5/3} m_1 m_2 M^{-1/3} \\ \dot{\omega} &= 3 \left(\frac{P_b}{2\pi}\right)^{-5/3} (T_\odot M)^{2/3} (1-e^2)^{-1}, \quad r = T_\odot m_2, \\ \gamma &= e \left(\frac{P_b}{2\pi}\right)^{1/3} T_\odot^{2/3} M^{-4/3} m_2 (m_1 + 2m_2), \quad s = x \left(\frac{P_b}{2\pi}\right)^{-2/3} T_\odot^{-1/3} M^{2/3} m_2^{-1}.\end{aligned}$$



Testing GR with GW observations: a **manifesto**

- **Consistency:** useful sanity checks, but hard to interpret statistically. P values are possible with much work. But would we ever believe an inconsistent result?
- **Parametric tests:** constraints on GR “constants” (PN coefficients, graviton mass) are useful proxies for increasing resolving power, but again hard to interpret. Apparent violations may focus our search for new physics.
- **Alternative theories:** new physics will be established by model comparison of GR with fully predictive alternative theories. It is a problem to establish Bayesian priors for alternative gravity, and for alternative-gravity parameters.

same as detection

other measurements, particle physics, astroparticles


GW150914 test of polarization



no discrimination
between S and T

$$\frac{P(s|S)}{P(s|T)} \sim 1$$


GW150914 test of polarization




no discrimination
between S and T

$$\frac{P(s|S)}{P(s|T)} \sim 1$$

Establishing alternative theories



Establishing alternative theories




new physics follows from establishing an **anomaly**:
we need to obtain convincing evidence that the data
prefers an alternative theory of gravity over GR

Testing GR with GW observations: a **manifesto**

- **Consistency:** useful sanity checks, hard to interpret statistically. P values are possible with much work. But would we ever believe an inconsistent result?
- **Parametric tests:** constraints on GR “constants” (PN coefficients, graviton mass) are useful proxies for increasing resolving power, but again hard to interpret. Apparent violations may focus our search for new physics.
- **Alternative theories:** new physics will be established by model comparison of GR with fully predictive alternative theories. It is a problem to establish Bayesian priors for alternative gravity, and for alternative-gravity parameters.
- **Size of effects:** detection SNR determines the magnitude of detectable waveform anomalies. 1% LIGO, up to 10^{-5} for LISA and future ground-based detectors.

How well can we hope to do? [MV, PRD 86, 2012]

- For a fixed false-alarm rate, we ask what **SNR** is needed to detect AG with 50% probability as a function of **fitting factor FF**, using the Bayesian odds ratio as “detection” statistic.




Testing GR with GW observations: a **manifesto**


- **Consistency:** useful sanity checks, hard to interpret statistically. P values are possible with much work. But would we ever believe an inconsistent result?
- **Parametric tests:** constraints on GR “constants” (PN coefficients, graviton mass) are useful proxies for increasing resolving power, but again hard to interpret. Apparent violations may focus our search for new physics.
- **Alternative theories:** new physics will be established by model comparison of GR with fully predictive alternative theories. It is a problem to establish Bayesian priors for alternative gravity, and for alternative-gravity parameters.
- **Size of effects:** detection SNR determines the magnitude of detectable waveform anomalies: 1% LIGO, up to 10^{-5} for LISA and future ground-based detectors.
- **Systematics:** beyond statistical-significance arguments, we will need a solid chain of evidence before we claim GR is violated.

Control of systematics


fundamental bias (Yunes)




astrophysical bias




mismodeling



calibration



glitches



Testing GR with GW observations: recommendations

- **Consistency:** useful sanity checks, hard to interpret statistically. P values are possible with much work. But would we ever believe an inconsistent result?
- **Parametric tests:** constraints on GR “constants” (PN coefficients, graviton mass) are useful proxies for increasing resolving power, but again hard to interpret. Apparent violations may focus our search for new physics.
- **Alternative theories:** new physics will be established by model comparison of GR with fully predictive alternative theories. It is a problem to establish Bayesian priors for alternative gravity, and for alternative-gravity parameters.
- **Size of effects:** detection SNR determines the magnitude of detectable waveform anomalies: 1% LIGO, up to 10^{-5} for LISA and future ground-based detectors.
- **Systematics:** beyond statistical-significance arguments, we will need a solid chain of evidence before we claim GR is violated.

develop!

Testing GR with GW observations: recommendations

- **Consistency:** useful sanity checks, hard to interpret statistically. P values are possible with much work. But would we ever believe an inconsistent result?
- **Parametric tests:** constraints on GR “constants” (PN coefficients, graviton mass) are useful proxies for increasing resolving power, but again hard to interpret. Apparent violations may focus our search for new physics.
- **Alternative theories:** new physics will be established by model comparison of GR with fully predictive alternative theories. It is a problem to establish Bayesian priors for alternative gravity, and for alternative-gravity parameters.
- **Size of effects:** detection SNR determines the magnitude of detectable waveform anomalies: 1% LIGO, up to 10^{-5} for LISA and future ground-based detectors.
- **Systematics:** beyond statistical-significance arguments, we will need a solid chain of evidence before we claim GR is violated.

develop!

fund!

Testing GR with GW observations: recommendations

- **Consistency:** useful sanity checks, hard to interpret statistically. P values are possible with much work. But would we ever believe an inconsistent result?
- **Parametric tests:** constraints on GR “constants” (PN coefficients, graviton mass) are useful proxies for increasing resolving power, but again hard to interpret. Apparent violations may focus our search for new physics.
- **Alternative theories:** new physics will be established by model comparison of GR with fully predictive alternative theories. It is a problem to establish Bayesian priors for alternative gravity, and for alternative-gravity parameters.
- **Size of effects:** detection SNR determines the magnitude of detectable waveform anomalies: 1% LIGO, up to 10^{-5} for LISA and future ground-based detectors.
- **Systematics:** beyond statistical-significance arguments, we will need a solid chain of evidence before we claim GR is violated.

develop!

fund!

control!

Testing GR with GW observations: recommendations

- **Consistency:** useful sanity checks, hard to interpret statistically. P values are possible with much work. But would we ever believe an inconsistent result?
- **Parametric tests:** constraints on GR “constants” (PN coefficients, graviton mass) are useful proxies for increasing resolving power, but again hard to interpret. Apparent violations may focus our search for new physics.
- **Alternative theories:** new physics will be established by model comparison of GR with fully predictive alternative theories. It is a problem to establish Bayesian priors for alternative gravity, and for alternative-gravity parameters.
- **Size of effects:** detection SNR determines the magnitude of detectable waveform anomalies: 1% LIGO, up to 10^{-5} for LISA and future ground-based detectors.
- **Systematics:** beyond statistical-significance arguments, we will need a solid chain of evidence before we claim GR is violated.

fold!

develop!

fund!

control!